

Hypoxia and Macrophages Polarization Defecets via Aryl Hydrocarbon Receptor (AHR) Perturbation

Saeed Mohammadi^{1,2*}

1. Stem Cell Research Center, Golestan University of Medical Sciences, Gorgan, Iran
2. Infectious Diseases Research Center, Golestan University of Medical Sciences, Gorgan, Iran

Article Type:

Letter to Editor

Article History:

Received: 5 Nov 2019

Revised: 15 Nov 2019

Accepted: 19 Nov 2019

*Correspondence:

Saeed Mohammadi,
Stem Cell Research Center, Golestan
University of Medical Sciences,
Gorgan, Iran.
saeed.mohammadi6578@yahoo.com



Abstract

Aryl hydrocarbon receptor (AHR) signaling has been introduced to be involved in macrophages polarization. The perturbation of AHR has been reported in hypoxia and related disorders. Here, I would like to highlight the significance of AHR in hypoxia-mediated macrophages polarization and suggest conducting further experiments on related subjects.

Keywords: Aryl Hydrocarbon Receptor; Hypoxia; Macrophages; Polarization

Statement

Macrophages are particular mononuclear phagocytic cells of the innate immune system, along with the monocytes and dendritic cells and are responsible for the regulation of immune responses and modulation of inflammation (1). Macrophages and their precursors are recognized for their significant phenotypic plasticity and their distinct functional roles which may be assumed to be contradictory; such as the secretion of both

pro- and anti-inflammatory cytokines, executing immunogenic and tolerogenic functions, and/or promoting both tissue repair and its injury (2). In response to (micro)environmental stimuli such as pathogens and their metabolites, or the apoptotic or dying cells, macrophages exhibit their functional and phenotypic heterogeneity (2). Accordingly, macrophages will be polarized into classically M1 activated (pro-inflammatory) and/or alternatively M2 activated (anti-inflammatory) subtypes (3).

Copyright© 2018, **Jorjani Biomedicine Journal** has published this work as an open access article under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial uses of the work while it is properly cited.



The imbalanced status of polarization in macrophages could be distinctive of some pathological conditions, such as autoimmune disorders (3), malignancies(4) and uncontrolled infectious diseases (5). Several molecules and signaling pathways have been proposed to be involved in macrophage polarization including peroxisome proliferator-activated receptor gamma (PPAR-gamma) (2), histone deacetylases (HDACs) (3), NF- κ B (5), interferon regulatory factors (IRFs) (5) and etc.

Recently, it has been suggested that aryl hydrocarbon receptor (AHR) signaling could be involved in macrophages polarization. In a recent publication by our team, we revealed that I3C-mediated activation of AHR could be associated with the overexpression of M2 and downregulation of M1 markers (enhancing M2 polarization), and exerting immunoregulatory effects on macrophages of SLE patients (6). Similarly, Shinde et al. showed that the activation of AHR by apoptotic/dying cell phagocytes maintains peripheral tolerance and is linked to the polarization of macrophages (7).

AHR is a cytoplasmic receptor and a member of the basic helix–loop–helix–(bHLH) superfamily of atranscription factors, which is normally responsible for metabolizing xenobiotics and is associated with several cellular process and immune responses (7). AHR is extremely responsive to a vast number of environmental and endogenous ligands (6). Hypoxia is a process in which cells are deprived from sufficient oxygen. It is an important determinant in several pathological processes of tumors, autoimmune disorders and fibrotic diseases (8). The imbalance polarization of macrophages is an important feature in most of the abovementioned disorders. It has been reported that hypoxia

could interfere with AHR binding to the target gene sequences and that AHR target gene expression could be perturbed in hypoxic environments (9). Interestingly, it is revealed that hypoxia can modify the polarization of macrophages and regulate the inflammatory microenvironment (10). Accordingly, I would like to highlight the significance of AHR signaling in hypoxia-mediated macrophages polarization. Then, AHR could be employed as a helpful target in modulation of inflammation in related disorders such as cancer and autoimmunity and considered as an interesting subject for future research.

References

1. Saghaeian-Jazi M, Mohammadi S, Sedighi S. Culture and differentiation of monocyte derived macrophages using human serum: an optimized method. *Zahedan J Res Med Sci.* 2016;18(6):e7362. <https://doi.org/10.17795/zjrms-7362>
2. Mohammadi S, Saghaeian-Jazi M, Sedighi S, Memarian A. Immunomodulation in systemic lupus erythematosus: induction of M2 population in monocyte-derived macrophages by pioglitazone. *Lupus.* 2017;26(12):1318-27. <https://doi.org/10.1007/s10067-017-3922-0>
3. Mohammadi S, Saghaeian-Jazi M, Sedighi S, Memarian A. Sodium valproate modulates immune response by alternative activation of monocyte-derived macrophages in systemic lupus erythematosus. *Clinical rheumatology.* 2018;37(3):719-27. <https://doi.org/10.1007/s10067-017-3922-0>
4. Pantano F, Berti P, Guida FM, Perrone G, Vincenzi B, Amato MMC, et al. The role of macrophages polarization in predicting prognosis of radically resected gastric cancer patients. *Journal of cellular and molecular medicine.* 2013;17(11):1415-21. <https://doi.org/10.1111/jcmm.12109>

5. Hoppstädter J, Hachenthal N, Valbuena-Perez JV, Lampe S, Astanina K, Kunze MM, et al. Induction of glucocorticoid-induced leucine zipper (GILZ) contributes to anti-inflammatory effects of the natural product curcumin in macrophages. *Journal of Biological Chemistry*. 2016;291(44):22949-60. <https://doi.org/10.1074/jbc.M116.733253>
6. Mohammadi S, Memarian A, Sedighi S, Behnampour N, Yazdani Y. Immunoregulatory effects of indole-3-carbinol on monocyte-derived macrophages in systemic lupus erythematosus: a crucial role for aryl hydrocarbon receptor. *Autoimmunity*. 2018;51(5):199-209. <https://doi.org/10.1080/08916934.2018.1494161>
7. Shinde R, Hezaveh K, Halaby MJ, Kloetgen A, Chakravarthy A, da Silva Medina T, et al. Apoptotic cell-induced AhR activity is required for immunological tolerance and suppression of systemic lupus erythematosus in mice and humans. *Nature immunology*. 2018;19(6):571-82. <https://doi.org/10.1038/s41590-018-0107-1>
8. Görlach A, Dimova EY, Petry A, Martínez-Ruiz A, Hernansanz-Agustín P, Rolo AP, et al. Reactive oxygen species, nutrition, hypoxia and diseases: problems solved? *Redox biology*. 2015;6:372-85. <https://doi.org/10.1016/j.redox.2015.08.016>
9. Vorrink SU, Severson PL, Kulak MV, Futscher BW, Domann FE. Hypoxia perturbs aryl hydrocarbon receptor signaling and CYP1A1 expression induced by PCB 126 in human skin and liver-derived cell lines. *Toxicology and applied pharmacology*. 2014;274(3):408-16. <https://doi.org/10.1016/j.taap.2013.12.002>
10. Ke X, Chen C, Song Y, Cai Q, Li J, Tang Y, et al. Hypoxia modifies the polarization of macrophages and their inflammatory microenvironment, and inhibits malignant behavior in cancer cells. *Oncology Letters*. 2019;18(6):5871-8. <https://doi.org/10.3892/ol.2019.10956>

How to cite:

Mohammadi M. Title: Hypoxia and Macrophages Polarization Defecets via Aryl Hydrocarbon Receptor (AHR) Perturbation. *Jorjani Biomedicine Journal*. 2019; 7(4): 1-3.